

On local symbolic approximation and resolution of ODEs using Implicit Function Theorem

Gianluca Argentini

gianluca.argentini@riellogroup.com

Research & Development Department

Riello Burners, 37048 San Pietro di Legnago (Verona), Italy

January 2006

Abstract

In this work the implicit function theorem is used for searching local symbolic resolution of differential equations. General results of existence for first order equations are proven and some examples, one relative to cavitation in a fluid, are developed. These examples seem to show that local approximation of non linear differential equations can give useful informations about symbolic form of possible solutions, and in the case a global solution is known, locally the accuracy of approximation can be good.

1 Introduction

The use of *Implicit Function* or *Dini's* Theorem for local resolution of differential equations is not new (see [7] for an historical example of application due to Bernoulli, and [9] for an application of theoretical existence for a non linear system of PDEs). But in the technical literature this method is very little widespread and limited to some specific situations. Moreover, for some physical and engineering aims, often usual numerical resolution of differential equations is not very useful, because the knowledge of an analytical dependence from physical or geometrical parameters is required (see e.g [3]) for the problem of searching analytical dependence for resistance coefficients in flows).

In this work some statements about local symbolic resolution of ordinary differential equations are established, and some examples, two very simple but illustrative the technique, and other two more technical and useful, are developed.

2 ODEs and Dini's Theorem

Let $[a, b]$, $a < b$, an interval of $\mathbb{R}$. If we denote with $y = y(x)$ a $C^n([a, b])$ function, for some n integer, $n > 0$, defined on $[a, b]$ and with $y^{(k)} = y^{(k)}(x)$ its k -th derivative, an ordinary differential equation (ODE) is an expression like this:

$$F(x, y(x), y^{(1)}(x), \dots, y^{(n)}(x)) = 0 \quad \forall x \in [a, b] \quad (1)$$

where F is a function $F : [a, b] \times \mathbb{R}^{n+1} \rightarrow \mathbb{R}$. Let $\partial_x F$ the partial derivative of F respect the x variable in $[a, b]$ and $\partial_k F$, $0 \leq k \leq n$, the partial derivative respect a suitable variable of $\mathbb{R}^{n+1}$ domain.

Suppose now $F \in C^m([a, b] \times \mathbb{R}^{n+1})$ for some integer $m > 0$.

If $T_0 = (x_0^0, p_0^0, \dots, p_n^0) \in [a, b] \times \mathbb{R}^{n+1}$ is a point such that $F(T_0) = 0$, and $\partial_k F(T_0) \neq 0$ for a particular value of k , then from *Dini's* or *Implicit Function* Theorem there are a constant $\delta > 0$, a costant $M > 0$, a neighbour U_0 of $(p_0^0, \dots, p_n^0)$ and a unique function $\phi : [a, b] \times \mathbb{R}^n \rightarrow \mathbb{R}$, $\phi \in C^m([a, b])$, such that

$$\begin{aligned} F(x, p_0, \dots, p_{k-1}, \phi(x, p_0, \dots, p_{k-1}, p_{k+1}, \dots, p_n), p_{k+1}, \dots, p_n) &= 0 \\ \forall (x, p_0, \dots, p_{k-1}, p_{k+1}, \dots, p_n) &\in (x_0 - \delta, x_0 + \delta) \times U_0 \end{aligned} \quad (2)$$

$$\phi(x_0^0, p_0^0, \dots, p_{k-1}^0, p_{k+1}^0, \dots, p_n^0) = p_k^0 \quad (3)$$

$$|\phi(x_0, p_0, \dots, p_{k-1}, p_{k+1}, \dots, p_n) - p_k^0| \leq M \quad (4)$$

and

$$\frac{\partial \phi}{\partial p_i} = - \frac{\partial_i F}{\partial_k F} \quad (5)$$

For a complete proof, see e.g. [4].

Let we suppose $n = 1$, so the differential equation is $F(x, y(x), y'(x)) = 0$. If the equation has order greater then 1, it is possible to write it as a system of order 1 differential equations (see e.g. [8]), and Dini Theorem has a natural extension to systems of equations.

Theorem 1 *Let $F(x, y(x), y'(x)) = 0$ an ODE, with $F \in C^1([a, b] \times \mathbb{R}^2)$ a real function $F(x, p, q)$, and let (x_0, p_0, q_0) such that $x_0 \in (a, b)$, $F(x_0, p_0, q_0) = 0$ and $\partial_q F(x_0, p_0, q_0) \neq 0$. Then exist $\delta > 0$, $\rho > 0$, a function $u \in C^1([x_0 - \delta, x_0 + \delta])$ and a function $\phi \in C^0([x_0 - \delta, x_0 + \delta] \times [p_0 - \rho, p_0 + \rho])$ such that $\phi(x_0, p_0) = q_0$ and*

$$u'(x) = \phi(x, u(x)), \quad F(x, u(x), \phi(x, u(x))) = 0 \quad \forall x \in [x_0 - \delta, x_0 + \delta] \quad (6)$$

Proof. From Dini Theorem exist $\delta_1 > 0$, $\rho_1 > 0$ and a function $\phi \in C^0([x_0 - \delta_1, x_0 + \delta_1] \times [p_0 - \rho_1, p_0 + \rho_1])$ such that

$$F(x, p, \phi(x, p)) = 0 \quad \forall (x, p) \in [x_0 - \delta_1, x_0 + \delta_1] \times [p_0 - \rho_1, p_0 + \rho_1] \quad (7)$$

We can consider the *differential problem*

$$y'(x) = \phi(x, y(x)), \quad y(x_0) = p_0 \quad (8)$$

Being ϕ a continuous function, from the theorem of *Peano* about existence of solutions for first order ordinary differential problems (or by local existence theorems) (see [2],[8]), exist δ_2 with $0 < \delta_2 \leq \delta_1$ and a function $u \in C^1([x_0 - \delta_2, x_0 + \delta_2])$ (local) solution of (8). Being u a continuous function and $u(x_0) = p_0$, exists a $0 < \delta_3 \leq \delta_2$ such that $u(x) \in [p_0 - \rho_1, p_0 + \rho_1] \quad \forall x \in [x_0 - \delta_3, x_0 + \delta_3]$. Then, from (7), if $x \in [x_0 - \delta_3, x_0 + \delta_3] \subseteq [x_0 - \delta_1, x_0 + \delta_1]$, we have $F(x, u(x), \phi(x, u(x))) = 0$, so with the choice $\delta = \delta_3$ and $\rho = \rho_1$ the proof is completed. $\square$

Therefore, under suitable hypothesis, ODE $F(x, y(x), y'(x)) = 0$ has at least local solutions. But the theorem says too that the original ODE can be *locally* written in a *new form*, i.e. in the expression $y'(x) = \phi(x, y(x))$, that is the *normal* form. In this sense, the term locally means that there is a neighbour of (x_0, p_0, q_0) where the original ODE can be written in the new form, and a local solution u exists such that $u(x_0) = p_0$ and $u'(x_0) = q_0$.

Example 1. Consider the ODE

$$2y(x) - y'(x) = 0 \quad (9)$$

in the interval $[a, b]$ where $a < 0$ and $b > 0$. The function F is $F(x, p, q) = 2p - q$, with $F \in C^\infty([a, b] \times \mathbb{R}^2)$ and $\partial_q F = -1$. Let $x_0 = 0$ and $p_0 = 1$; then $q_0 = 2$. From theorem 1 exist two functions u and ϕ such that $\phi(0, 1) = 2$ and $2p - \phi(x, p) = 0$ in a neighbour of $(0, 1)$. Then $\phi(x, p) = 2p$, so from (6) $u'(x) = 2u(x)$, that is $u(x) = ce^{2x}$. The constant c is determined from the condition $u(0) = p_0 = 1$, hence $c = 1$.

In the case $\partial_q F(x_0, p_0, q_0) = 0$, we can consider the possibility of a local expression like $y(x) = \psi(x, y'(x))$, as stated in the following

Theorem 2 Let $F(x, y(x), y'(x)) = 0$ an ODE, with $F \in C^1([a, b] \times \mathbb{R}^2)$ a real function $F(x, p, q)$, and let (x_0, p_0, q_0) such that $x_0 \in (a, b)$, $F(x_0, p_0, q_0) = 0$ and $\partial_p F(x_0, p_0, q_0) \neq 0$. Then exist $\delta > 0$, $\rho > 0$ and a function $\psi \in C^0([x_0 - \delta, x_0 + \delta] \times [q_0 - \rho, q_0 + \rho])$ such that $\psi(x_0, q_0) = p_0$ and

$$F(x, \psi(x, q), q) = 0 \quad \forall (x, q) \in [x_0 - \delta, x_0 + \delta] \times [q_0 - \rho, q_0 + \rho] \quad (10)$$

If exists a solution $u \in C^1([x_0 - \sigma, x_0 + \sigma])$, with $0 < \sigma < \delta$, for the problem

$$\begin{cases} y(x) = \psi(x, y'(x)) & x \in [x_0 - \sigma, x_0 + \sigma] \\ y'(x_0) = q_0 \end{cases} \quad (11)$$

then

$$F(x, u(x), u'(x)) = 0 \quad \forall x \in [x_0 - \sigma, x_0 + \sigma] \quad (12)$$

Proof. The first statement is a direct consequence of Dini's Theorem. Let then u a solution of the problem (11), with $\sigma_1 < \delta$. From the fact that u' is continuous in the interval $[x_0 - \sigma_1, x_0 + \sigma_1]$, we can choose the constant σ so that $u'(x) \in [q_0 - \rho, q_0 + \rho]$ for every $x \in [x_0 - \sigma, x_0 + \sigma]$. Then, from (10) follows that $F(x, u(x), u'(x)) = 0$ for every $x \in [x_0 - \sigma, x_0 + \sigma]$. $\square$

Therefore, a local solution for the new differential equation $y(x) = \psi(x, y'(x))$ is a local solution for the original ODE $F(x, y(x), y'(x)) = 0$.

Example 2. Consider the non linear ODE

$$y'(x)^2 + y(x)^2 - 1 = 0 \quad (13)$$

The function F is $F(x, p, q) = q^2 + p^2 - 1$, with $\partial_p F = 2p$ and $\partial_q F = 2q$. If we consider the point $T_0 = (\frac{\pi}{2}, 1, 0)$, we have $\partial_q F(T_0) = 0$ and $\partial_p F(T_0) = 2$, so from the first statement of theorem 2, if $|q| \leq 1$ exists the function $\psi = \psi(x, q) = \sqrt{1 - q^2}$ such that (locally) $p = \sqrt{1 - q^2}$. Then we can consider the problem

$$\begin{cases} y(x) = \sqrt{1 - y'(x)^2} & \forall x \in [0, \pi] \\ y'(\frac{\pi}{2}) = 0 \end{cases} \quad (14)$$

The function $u(x) = \sin(x)$ is a solution of this problem, with $|u'(x)| \leq 1$ as required, and is a local (but in this case global too) solution of the original ODE. Note that finding the solution of the original equation implies the use of complex variables: the internal command `DSolve[y'[x]^2 + y[x]^2 - 1 == 0, y[x], x]` of the software *Mathematica* gives the integrals

$$u(x) = 1 - 2\sin^2 \left[\frac{1}{2}(x - ic) \right], \quad v(x) = 1 - 2\sin^2 \left[\frac{1}{2}(-x - ic) \right] \quad (15)$$

where i is the imaginary unit and c a complex constant. We obtain our previous local solution respectively for, e.g., $c = i\frac{3}{2}\pi$ and $c = -i\frac{3}{2}\pi$.

3 Local symbolic approximation for an ODE

From implicit function theorem, if F has continuous partial derivatives, using Taylor expansion near the point (x_0, p_0) (or (x_0, q_0)) and the chain rule for the derivatives of composite function it is possible to write the following expression for the function ϕ (or ψ):

$$\begin{aligned} \phi(x, p) &= \phi(x_0, p_0) + [\partial_x \phi(x_0, p_0)](x - x_0) + [\partial_p \phi(x_0, p_0)](p - p_0) + \\ &+ \left[\frac{1}{2} \partial_{xx}^2 \phi(x_0, p_0) \right] (x - x_0)^2 + \left[\frac{1}{2} \partial_{pp}^2 \phi(x_0, p_0) \right] (p - p_0)^2 + \dots \end{aligned} \quad (16)$$

with, e.g. for x -derivation,

$$\partial_x \phi(x_0, p_0) = -\frac{\partial_x F}{\partial_q F}(x_0, p_0, q_0) \quad (17)$$

$$\partial_{xx}^2 \phi(x_0, p_0) = -\frac{(\partial_x \phi)^2 \partial_{qq}^2 F + 2\partial_x \phi \partial_{xq}^2 F + \partial_{xx}^2 F}{\partial_q F} \quad (18)$$

where the derivatives of ϕ are evaluated at (x_0, p_0) and those of F at (x_0, p_0, q_0) . The expansion gives a local analytical approximation of the original ODE. If for this equation a global solution doesn't exist or it is not given by elementary functions, the expansion could be a more simple equation to solve. From previous theorems, a solution of this equation is an approximate local solution of the original ODE.

If $F \in C^1([a, b] \times \mathbb{R}^2)$, $F(x_0, p_0, q_0) = 0$ and $\partial_q F(x_0, p_0, q_0) \neq 0$, from previous identities the first order differential equation $F(x, y(x), y'(x))$ has the following local *approximation of first order*:

$$\begin{aligned} y'(x) = q_0 + \left[\frac{\partial_x F}{\partial_q F}(x_0, p_0, q_0) \right] x_0 + \left[\frac{\partial_p F}{\partial_q F}(x_0, p_0, q_0) \right] p_0 - \\ - \left[\frac{\partial_x F}{\partial_q F}(x_0, p_0, q_0) \right] x - \left[\frac{\partial_p F}{\partial_q F}(x_0, p_0, q_0) \right] y(x) \end{aligned} \quad (19)$$

If $\partial_p F(x_0, p_0, q_0) = 0$, the solution of this first order ODE is

$$y(x) = c + \left[q_0 + \frac{\partial_x F}{\partial_q F} x_0 \right] x - \frac{1}{2} \frac{\partial_x F}{\partial_q F} x^2 \quad (20)$$

otherwise its general solution is (see e.g [1])

$$y(x) = p_0 + \frac{\partial_x F \partial_q F}{[\partial_p F]^2} + \frac{\partial_x F x_0 + \partial_q F q_0}{\partial_p F} - \frac{\partial_x F}{\partial_p F} x + c e^{-\frac{\partial_p F}{\partial_q F} x} \quad (21)$$

where c is a constant and the partial derivatives are all calculated at point (x_0, p_0, q_0) . Imposing the condition $y(x_0) = p_0$ we obtain

$$c = -\frac{[\partial_x F + \partial_p F q_0] \partial_q F e^{\frac{\partial_p F}{\partial_q F} x_0}}{[\partial_p F]^2} \quad (22)$$

while, with this value of the constant c , we have $y'(x_0) = q_0$ as required.

So we have proven the following

Proposition 1 *In the hypothesis of Theorem 1, the ODE $F(x, y(x), y'(x)) = 0$ has a local solution, approximated to first order, of the form*

$$y(x) = a_0 + b_0 x + c_0 x^2 \quad \text{if } \partial_p F(x_0, p_0, q_0) = 0 \quad (23)$$

$$y(x) = a_0 + b_0 x + c_0 e^{d_0 x} \quad \text{if } \partial_p F(x_0, p_0, q_0) \neq 0 \quad (24)$$

where a_0 , b_0 , c_0 and d_0 are constants depending from (x_0, p_0, q_0) .

Example 1 bis. For the ODE $2y(x) - y'(x) = 0$, being $F(x, p, q) = 2p - q$, at point $(0, 1, 2)$ we obtain $\partial_x F = 0$, $\partial_p F = 2$ and $\partial_q F = -1$, so that $a_0 = b_0 = 0$, $c_0 = 1$ and $d_0 = 2$, therefore from Proposition 1 the local approximated solution of first order is $y(x) = e^{2x}$, which is the exact local (global) solution of the original equation, satisfying the condition $y(0) = 1$.

If $F \in C^1([a, b] \times \mathbb{R}^2)$, $F(x_0, p_0, q_0) = 0$ and $\partial_p F(x_0, p_0, q_0) \neq 0$, in a similar fashion the first order differential equation $F(x, y(x), y'(x)) = 0$ has the following local *approximation of first order*:

$$\begin{aligned} y(x) = p_0 + \left[\frac{\partial_x F}{\partial_p F}(x_0, p_0, q_0) \right] x_0 + \left[\frac{\partial_q F}{\partial_p F}(x_0, p_0, q_0) \right] q_0 - \\ - \left[\frac{\partial_x F}{\partial_p F}(x_0, p_0, q_0) \right] x - \left[\frac{\partial_q F}{\partial_p F}(x_0, p_0, q_0) \right] y'(x) \end{aligned} \quad (25)$$

If $\partial_q F(x_0, p_0, q_0) = 0$, the local approximated solution is simply

$$y(x) = p_0 + \frac{\partial_x F}{\partial_p F} x_0 - \frac{\partial_x F}{\partial_p F} x \quad (26)$$

while in the case $\partial_q F(x_0, p_0, q_0) \neq 0$ one can verify that the local solution is the same as (21). So the following statement holds:

Proposition 2 *In the hypothesis of Theorem 2, the ODE $F(x, y(x), y'(x)) = 0$ has a local solution, approximated to first order, of the form*

$$y(x) = a_0 + b_0 x \quad \text{if } \partial_q F(x_0, p_0, q_0) = 0 \quad (27)$$

$$y(x) = a_0 + b_0 x + c_0 e^{d_0 x} \quad \text{if } \partial_q F(x_0, p_0, q_0) \neq 0 \quad (28)$$

where a_0 , b_0 , c_0 and d_0 are constants depending from (x_0, p_0, q_0) .

Example 2 bis. For the ODE $y'(x)^2 + y(x)^2 - 1 = 0$, being $F(x, p, q) = p^2 + q^2 - 1$, at point $(\frac{\pi}{2}, 1, 0)$ we have $\partial_x F = 0$, $\partial_p F = 2$ and $\partial_q F = 0$, therefore $a_0 = 1$, $b_0 = 0$, $c_0 = 0$ and $d_0 = 2$. From previous proposition, a local approximated solution of first order is $y(x) = 1$. This result, compared with the exact solution $y(x) = \sin(x)$, is not satisfactory. A greater order of approximation is required.

If $F \in C^n([a, b] \times \mathbb{R}^2)$ for a suitable integer $n > 0$, $F(x_0, p_0, q_0) = 0$ and, e.g., $\partial_p F(x_0, p_0, q_0) \neq 0$, it is possible consider higher order derivatives for the implicit function ψ , as (18) or

$$\partial_{qq}^2 \psi(x_0, q_0) = - \frac{(\partial_q \psi)^2 \partial_{pp}^2 F + 2 \partial_q \psi \partial_{pq}^2 F + \partial_{qq}^2 F}{\partial_p F} \quad (29)$$

Then, from Taylor expansion of ψ , we can write the following approximation of order (1, 2) (1 for x -variable, 2 for q -variable) for an ODE:

$$\begin{aligned}
y(x) = & p_0 + \\
& + \frac{1}{2\partial_p F} \left[2\partial_x F x_0 + 2\partial_q F q_0 - \left(\frac{[\partial_q F]^2 \partial_{pp}^2 F}{[\partial_p F]^2} - 2\frac{\partial_q F}{\partial_p F} \partial_{pq}^2 F + \partial_{qq}^2 F \right) q_0^2 \right] - \\
& - \frac{1}{\partial_p F} \left[\partial_x F x + \left(\partial_q F q - \left(\frac{[\partial_q F]^2 \partial_{pp}^2 F}{[\partial_p F]^2} + 2\frac{\partial_q F}{\partial_p F} \partial_{pq}^2 F + \partial_{qq}^2 F \right) q_0 \right) y'(x) \right] \\
& - \frac{1}{2\partial_p F} \left(\frac{[\partial_q F]^2 \partial_{pp}^2 F}{[\partial_p F]^2} - 2\frac{\partial_q F}{\partial_p F} \partial_{pq}^2 F + \partial_{qq}^2 F \right) [y'(x)]^2
\end{aligned}$$

Expressions like this gives more accurate local solutions.

Example 2 ter. For the differential equation of Example 2 we have $\partial_x F(\frac{\pi}{2}, 1, 0) = \partial_q F(\frac{\pi}{2}, 1, 0) = \partial_{pq}^2 F(\frac{\pi}{2}, 1, 0) = 0$, $\partial_p F(\frac{\pi}{2}, 1, 0) = \partial_{pp}^2 F(\frac{\pi}{2}, 1, 0) = \partial_{qq}^2 F(\frac{\pi}{2}, 1, 0) = 2$, so from previous formula the approximation is

$$y(x) = 1 - \frac{1}{2}[y'(x)]^2 \quad (30)$$

The solution to this non linear equation, with condition $y(\frac{\pi}{2}) = 1$, is (computing it using `DSolve[{y[x]==1-1/2*y'[x]^2,y[Pi/2]==1},y[x],x]` of *Mathematica*)

$$u(x) = 1 - \frac{\pi^2}{8} + \frac{\pi}{2}x - \frac{1}{2}x^2 \quad (31)$$

This expression is the second order truncated Taylor expansion of $\sin(x)$ centered at $x = \frac{\pi}{2}$, so we have obtained a good approximation for the local solution found in Example 2. Note that, being in this case $\partial_x F = \partial_{xx}^2 F = 0$, from (17) and (18) follows that $\partial_{xx}^2 \psi(\frac{\pi}{2}, 0) = 0$, hence second order derivative respect x -variable is not useful for a more accurate expression of the local solution.

4 Relationship with series expansion of solution

The last example seems establish a relation between symbolic expression of local solution found using implicit function technique and powers expression of solution found using series expansion of the unknown function in the original ODE. Let us consider the following symbolic expansion of second order, centered at $x_0 = \frac{\pi}{2}$, for the first member of the original equation $y(x)^2 + y'(x)^2 - 1 = 0$ (using the *Mathematica* command `ode = Series[y[x]^2 + (D[y[x],x])^2 - 1, {x, Pi/2, 2}]`, see [5]):

$$\begin{aligned}
& -1 + y\left(\frac{\pi}{2}\right)^2 + y'\left(\frac{\pi}{2}\right)^2 - y'\left(\frac{\pi}{2}\right) \left[y\left(\frac{\pi}{2}\right) + y''\left(\frac{\pi}{2}\right) \right] (\pi - 2x) + \\
& \left[y'\left(\frac{\pi}{2}\right)^2 + y''\left(\frac{\pi}{2}\right) \left(y\left(\frac{\pi}{2}\right) + y''\left(\frac{\pi}{2}\right) \right) + y'\left(\frac{\pi}{2}\right) y'''\left(\frac{\pi}{2}\right) \right] \left(-\frac{\pi}{2} + x \right)^2
\end{aligned}$$

Then we solve the algebraic system formed by the expanded differential equation and the two conditions $y(\frac{\pi}{2}) = 1$, $y'(\frac{\pi}{2}) = 0$ (command `Solve[{ode == 0, y[Pi/2] == 1, (D[y[x], x] /. x -> Pi/2) == 0}]`). The result consists of two possible choices: $y''(\frac{\pi}{2}) = 0$ and $y''(\frac{\pi}{2}) = -1$. The expansion of the solution is $y(x) = y(\frac{\pi}{2}) + y'(\frac{\pi}{2})(x - \frac{\pi}{2}) + \frac{1}{2}y''(\frac{\pi}{2})(x - \frac{\pi}{2})^2$; for $y''(\frac{\pi}{2}) = 0$ we obtain the trivial solution $y = 1$, for $y''(\frac{\pi}{2}) = -1$ the solution is $y(x) = 1 - \frac{\pi^2}{8} + \frac{\pi}{2}x - \frac{1}{2}x^2$. Therefore the series expansion method is computationally equivalent to the method of implicit function, but not logically: the first is based on a direct expansion of the unknown function in the original differential equation, while the second is based on an approximation of this ODE for finding a symbolic solution of the new equation.

5 Finer examples

Example 3. Let us consider the non linear ODE (see [8])

$$y'(x) = -3\sin(x) [y(x)]^{\frac{4}{3}} \quad (32)$$

This equation is integrable by elementary functions:

$$y(x) = -\frac{27}{[c + 3\cos(x)]^3} \quad (33)$$

The associated function F is $F(x, p, q) = -3\sin(x)p^{\frac{4}{3}} - q$, and consider the point $T_0 = (\frac{\pi}{3}, 1, -3\sqrt[3]{3})$ where F is null. The following identities hold:

$$\partial_x F = -3\cos(x)p^{\frac{4}{3}}, \quad \partial_p F = -4\sin(x)p^{\frac{1}{3}}, \quad \partial_q F(T_0) = -1 \quad (34)$$

$$\partial_x F(T_0) = -\frac{3}{2}, \quad \partial_p F(T_0) = -2\sqrt{3}, \quad \partial_q F(T_0) = -1 \quad (35)$$

We want a Taylor expansion for the implicit function $q = \phi(x, p)$ of order $(3, 1)$, that is of third order respect to x -variable and first order respect to p -variable. We use expressions (17) and (18) for ϕ , and the following formula for the third derivative:

$$\begin{aligned}
\partial_{xxx}^3 \phi(x_0, p_0) = & -\frac{\left[3\partial_x \phi \partial_{xx}^2 \phi \partial_{pp}^2 F + [\partial_x \phi]^3 \partial_{ppp}^3 F + 3\partial_{xx}^2 \phi \partial_{xp}^2 F \right]}{\partial_q F} - \\
& -\frac{\left[3[\partial_x \phi]^2 \partial_{xpp}^3 F + 3\partial_x \phi \partial_{xpp}^3 F + \partial_{xxx}^3 F \right]}{\partial_q F}
\end{aligned} \quad (36)$$

Computation gives the following values for the derivatives at T_0 :

$$\partial_x \phi = -\frac{3}{2}, \quad \partial_{xx}^2 \phi = 3\frac{\sqrt{3}}{2}, \quad \partial_{xxx}^3 \phi = 3\frac{\sqrt{3}}{2}, \quad \partial_p \phi = -2\sqrt{3} \quad (37)$$

so the new approximated ODE is

$$y'(x) = -3\frac{\sqrt{3}}{2} - \frac{3}{2}\left(x - \frac{\pi}{3}\right) + \frac{3}{4}\sqrt{3}\left(x - \frac{\pi}{3}\right)^2 + \frac{\sqrt{3}}{4}\left(x - \frac{\pi}{3}\right)^3 - 2\sqrt{3}(y(x) - 1) \quad (38)$$

Solving this equation with conditions $y(\frac{\pi}{3}) = 1$, $y'(\frac{\pi}{3}) = -3\frac{\sqrt{3}}{2}$ (e.g with command **DSolve**) one obtains a solution of the form $y(x) = e^{ax}(b + cx + dx^2 + ex^3)$ which Taylor expansion of third order centered at $\frac{\pi}{3}$ is

$$y(x) = 1 + \frac{\sqrt{3}}{2}(\pi - 3x) + \frac{5}{12}(\pi - 3x)^2 + \frac{1}{4\sqrt{3}}(\pi - 3x)^3 \quad (39)$$

The exact solution of the original differential equation is $y(x) = -\frac{27}{[-\frac{9}{2} + 3\cos(x)]^3}$ which Taylor expansion of third order centered at $\frac{\pi}{3}$ is

$$y(x) = 1 + \frac{\sqrt{3}}{2}(\pi - 3x) + \frac{5}{12}(\pi - 3x)^2 + \frac{2}{9\sqrt{3}}(\pi - 3x)^3 \quad (40)$$

Therefore the implicit function method gives a very good approximated symbolic solution.




Figure 1: Comparison between exact and approximated solution in the interval $[\frac{\pi}{3} - 1, \frac{\pi}{3} + 1]$; the exact solution is dashed.

Example 4. Consider a spherical bubble of a compressible gas immersed in an incompressible nonviscous fluid. If this fluid is flowing in a vane of a mechanical structure as rotating pump, presence of bubbles (*cavitation*) can give increment of noise and reduction of efficiency. Hence it is important to estimate the time t_c necessary for the collapse of the bubble, assuming that gas pressure is negligible compared to external fluid pressure p_f . If $y(t)$ is the radius of the bubble, it is possible to derive from Navier-Stokes equations the following ODE (see [6])

$$\frac{2}{3}y(t)y''(t) + y'(t)^2 = -\frac{2}{3}\frac{p_f}{\rho} \quad (41)$$

It is convenient to express this ODE as a first order equation, which is equivalent if one doesn't consider constant solutions $y(t) = k$ (see [6] for details):

$$y(t)^3 y'(t)^2 = \frac{2}{3}\frac{p_f}{\rho} [R_0^3 - y(t)^3] \quad (42)$$

where ρ is the density of the external fluid and R_0 is the radius of the bubble at time $t = 0$. Without considering the trivial solution $y(t) = R_0$, which is not physically admissible, this equation is not symbolically integrable using elementary functions.

For simplicity, but without lack of generality, suppose $\rho = 1$ and the pressures normalized with respect to the value of p_f , so that $p_f = 1$. Let then $F(t, p, q) = \frac{2}{3}R_0^3 - (\frac{2}{3} + q^2)p^3$ the associated function F defining the implicit solution. From $F(0, R_0, q_0) = 0$ follows $q_0 = 0$, as required from the physics of the problem, therefore the point (t_0, p_0, q_0) is $T_0 = (0, R_0, 0)$. The following formulas hold for first and second derivatives of F :

$$\begin{aligned} \partial_t F &= 0, & \partial_p F &= -(2 + 3q^2)p^2, & \partial_q F &= -2p^3q, \\ \partial_{pp}^2 F &= -(4 + 6q^2)p, & \partial_{qq}^2 F &= -2p^3, & \partial_{pq}^2 F &= -6p^2q \end{aligned}$$

from which we have $\partial_p F(T_0) = -2R_0^2$, $\partial_q F(T_0) = \partial_{pq}^2 F(T_0) = 0$, $\partial_{pp}^2 F(T_0) = -4R_0$ and $\partial_{qq}^2 F(T_0) = -2R_0^3$. Then, using (16), (17) and (29), we can use the following approximation of order (1,2) (1 for y , 2 for y'), centered at $(0, R_0, 0)$, for (42):

$$y(t) = R_0 \left[1 - \frac{1}{2}y'(t)^2 \right] \quad (43)$$

This equation is a non linear ODE, but it is integrable by elementary functions. Its particular solution with initial condition $y(t) = R_0$ is (e.g using command `DSolve[{y[t]==R_0(1-1/2 y'[t]^2), y[0]==R_0}, y[t], t]`)

$$y(t) = -\frac{1}{2R_0}t^2 + R_0 \quad (44)$$

from which we have our estimate for the time t_c of collapse:

$$t_c = \sqrt{2}R_0 \quad (45)$$

From this formula one can observe the importance of the initial radius of the bubble and consequently develop suitable actions. It is interesting to compare this result with the estimate from a numerical resolution of the original ode (41) (`NDSolve[{2/3 y[t] y'[t] + y'[t]^2 == -2/3, y[0] == R_0}, y[t], {t, 0, 0.12}]`).




Figure 2: Comparison between symbolic approximated solution of the reduce ODE (continuous line) and numerical solution of the original ODE (dashed line); case $p_f = 1$, $\rho = 1$, $R_0 = 0.1$; time is in horizontal axis, radius in vertical axis.

The figure 2 shows that symbolic solution of the reduce ODE is a good approximation of the implicit solution of the original equation.

6 Technical notes

We have use internal commands `DSolve` and `NDSolve` of *Mathematica* ver. 5.2 for symbolic and numerical resolution of ODEs cited in the examples. Integrability of these equations using elementary functions has been tested using `DSolve` (see [5]).

References

- [1] Boyce, W.E. and DiPrima, R.C., *Elementary Differential Equations*, 6th edition, Wiley, (1996)
- [2] Birkhoff, G. and Rota G., *Ordinary Differential Equation*, 4th edition, Wiley, New York, (1989)
- [3] Buttazzo, G. and Kawohl, B., *On Newton's Problem of Minimal Resistance*, in Wilson, R. and Gray, J., *Mathematical Conversations*, Springer-Verlag, New York, (2001)
- [4] Courant, R. *Differential and Integral Calculus*, Vol. 2, Wiley, (1988)

- [5] Coombes, K.; Hunt, B.; Lipsman, R.; Osborn, J. and Stuck, G. *Differential Equations with Mathematica*, 2nd edition, John Wiley & Sons, Inc. (1998)
- [6] Malek-Madani, R. *Advanced Engineering Mathematics*, Addison-Wesley, (1998)
- [7] Mingari, G. and Ritelli, D. *An Historical Outline of the Theorem of Implicit Function*, Divulgaciones Matematicas **10**(2), 171-180, (2002)
- [8] Sanchez, D. *Ordinary Differential Equations and Stability Theory: an introduction*, Dover Publications, New York, (1979)
- [9] Valent, T. *Boundary value problems of finite elasticity: local theorems on existence, uniqueness, and analytic dependence on data*, Springer, New York, (1988)